

***BENZOYLALKYLINDOLEPYRIDINIUM COMPOUNDS
AND PHARMACEUTICAL COMPOSITIONS COMPRISING SUCH COMPOUNDS***

The present application claims priority from U.S. Provisional Application No.
5 60/256,556, filed on December 18, 2000.

FIELD

The present invention concerns benzoylalkylindolepyridinium compounds,
pharmaceutical compositions comprising such compounds, and methods for making and using
10 such compounds and compositions.

BACKGROUND

Viruses cause a variety of human and animal illnesses. Many are relatively harmless
and self-limiting, but the other end of the spectrum includes acute life-threatening illnesses such
15 as hemorrhagic fever, and prolonged serious illnesses such as hepatitis B and acquired immune
deficiency syndrome (AIDS). Unlike bacterial infections, where numerous suitable antibiotic
drugs are usually available, there are relatively few effective antiviral treatments.

A. Viruses

20 Viruses consist of a nucleic acid surrounded by one or more proteins. A virus's nucleic
acid typically comprises relatively few genes, embodied either as DNA or RNA. DNA
genomes may be single or double-stranded (examples include hepatitis B virus and herpes
virus). RNA genomes may be single strand sense (so-called positive-strand genomes; examples
include poliovirus), single strand or segmented antisense (so-called negative-strand genomes;
25 examples include HIV and influenza virus), or double-stranded segmented RNA genomes
(examples include rotavirus, an acute intestinal virus).

Retroviruses represent a particular family of negative stranded RNA virus. The term
"retrovirus" means that in the host cell the viral RNA genome is transcribed into DNA. Thus,
information is not passing in the "normal" direction, from DNA to RNA to proteins, but rather
30 in a "retrograde" direction, from RNA to DNA. To accomplish this change in direction, a
retrovirus has one of a unique class of enzymes referred to as the reverse transcriptases. These
enzymes are RNA-dependent DNA polymerases -- that is, they synthesize DNA strands using

the viral RNA genome as a template. Each species of retrovirus has its own reverse transcriptase. Once the reverse transcriptase copies the retroviral RNA genome, it uses its inherent *DNA-dependent* DNA polymerase activity—that is, the ability to synthesize DNA copied from other DNA—to generate a double-stranded DNA version of the viral DNA
5 genome.

HIVs (human immunodeficiency viruses) are retroviruses of the lentivirus subfamily. The two known subfamily members that infect humans are called HIV-1 and HIV-2 (simian immunodeficiency virus, or SIV, is a closely related lentivirus that infects monkeys). Once the virus gains entry into the body, it attaches to human immune cells that express the CD4 receptor
10 on their surface (CD4+ cells). CD4+ cells (which include “helper” and lymphocytes and monocytes), become the primary repository for the virus. HIV-1 isolates are categorized into two broad groups, group M and group O. Group O comprises eight subtypes or clades, designated A through H.

15 **B. Viral Therapeutics**

Currently, only a limited number of drugs are approved for treating viral infections, such as human immune deficiency virus Type 1 (HIV-1) infection. Two broad families of anti-HIV drugs include the viral protease inhibitors, and the reverse transcriptase (RT) inhibitors. There are three main classes of RT inhibitors: (1) dideoxynucleoside (ddN) analogs, (2) acyclic
20 nucleoside phosphonate (ANP) analogs, and (3) non-nucleoside reverse transcriptase inhibitors (NNRTIs).


The ddN and ANP nucleoside analog drugs are phosphorylated inside the cell. Once phosphorylated, they bind to the RT’s substrate binding site. This is the site where the RT binds nucleotides (dATP, dCTP, dGTP, or dTTP, collectively referred to as dNTPs) so that they can
25 be added to the growing DNA chain. When a nucleoside analog drug binds to the RT substrate binding site, it is integrated into the DNA, just as a normal dNTP would. But the enzyme cannot subsequently add dNTPs onto the incorporated nucleoside analog. Thus, the two classes of nucleoside analogs function as “chain terminators,” and thereby limit HIV replication. These drugs have proven clinically effective against HIV infection, but resistance rapidly emerges due
30 to mutations in and around the RT active site.

NNRTIs do not require phosphorylation or function as chain terminators, and do not bind at the substrate (dNTP) binding site. Known NNRTIs bind to a specific region outside the

RT active site, and cause conformational changes in the enzyme that render it inactive. Known NNRTIs are highly potent and relatively non-toxic agents that are extremely selective for inhibition of HIV-1 RT. However, like the nucleoside analogs, their use is limited by the rapid emergence of resistant strains. In addition, they do not inhibit the RT activity of HIV-2, SIV
5 and possibly some HIV-1 Group O isolates, nor do they prevent these viruses from replicating.

C. *Pyrido-Indole Compounds*

Ryabova et al. describe certain pyrido-indole compounds in "2-Formyl-3-Aryl-aminoindoles in the Synthesis of 1,2- and 1,4-Dehydro-5H-Pyrido-[3,2-b]-Indole (δ carboline) Derivatives," *Pharmaceutical Chemistry Journal*, 30:579-583 (1996). For example, Ryabova et
10 al. describe 1-(4-nitrophenyl)-2-dimethylamino-3-cyano-4-(2-oxo-propyl)-5-methyl-1,4-dehydro-5H-pyrido [3,2-b]-indole (Compound 2).



Compound 2

15 No biological data is provided for this compound.

D. *Conclusion*


The treatment of viral diseases, such as HIV disease, has been significantly advanced by the recognition that combining different drugs with specific activities against different
20 biochemical functions of the virus can help reduce the rapid development of drug resistant viruses. However, even with combined treatments, multi-drug resistant strains of the virus have

emerged. Therefore, there is a continuing need to develop new drugs, particularly antiviral drugs that act specifically at different steps of the viral infection and replication cycle.

SUMMARY

- 5 The disclosed invention provides new antiviral compounds and pharmaceutical compositions comprising such compounds, particularly antiretroviral compounds and compositions, that address many of the problems noted above. These compounds, referred to as benzoylalkylindolepyridinium compounds (BAIPs), are effective against HIV isolates that have developed mutations rendering conventional drugs ineffective in their treatment. The BAIPs
- 10 apparently do not require intracellular phosphorylation nor bind to the RT active site, which distinguishes their mechanism of action from the ddN and ANP nucleoside analog drugs. The BAIPs also may be distinguished from the NNRTIs, in part because the BAIPs bind to a different site on the RT enzyme. Moreover, unlike the NNRTIs, BAIPs of the present invention have been shown to be effective for limiting HIV-1, HIV-2, and SIV proliferation. Thus,
- 15 BAIPs are broadly antiviral, non-nucleoside reverse transcriptase inhibitors (BANNRTIs).


Novel BAIPs have Formula I below.




Formula I

With reference to Formula I, R is selected from the group consisting of hydrogen and lower aliphatic, particularly lower alkyl, such as methyl. The nitro group (-NO₂) can be at any ring position, i.e., *ortho*, *meta* or *para* to the ring nitrogen, but typically is in the *para* position.


One novel compound of the present invention is shown below (Compound 2).



The present invention also provides a method for treating a subject, such as treating viral infections. The method comprises providing a compound having Formula II.

**Formula II**

With reference to Formula II, R₁ is selected from the group consisting of hydrogen and lower aliphatic, particularly lower alkyl, such as methyl; and R₂ is selected from the group consisting of -CH₂COCH₃ and



where R₁ is as stated for Formula II.

The compound is administered in effective amounts to subjects, such as a human or simian. A person of ordinary skill in the art will realize that the effective amount can vary. However, solely by way of guidance, an effective amount typically is from about 0.1 mg/kg body weight per day, to about 200 mg/kg body weight per day, in single or divided doses. The compound, or compounds, can be administered in any of a number of ways, including without limitation, topically, orally, intramuscularly, intranasally, subcutaneously, intraperitoneally, intravenously, or combinations thereof. The currently preferred administration method is

intravenous. Such compounds also can be administered as pharmaceutical compositions, and hence may include other materials commonly found in pharmaceutical preparations, including other therapeutic agents.

The present invention also provides compositions comprising amounts of a compound
5 or compounds effective to treat diseases, particularly viral infections. One likely mechanism of action is by inhibition of reverse transcriptase, and therefore effective amounts can be amounts sufficient to inhibit reverse transcriptase. Such compositions may further comprise inert carriers, excipients, diagnostics, direct compression binders, buffers, stabilizers, fillers, disintegrants, flavors, colors, lubricants, other active ingredients, other materials conventionally
10 used in the formulation of pharmaceutical compositions, and mixtures thereof.

A method for treating a subject, particularly mammals, such as humans and simians, also is provided. The method first comprises providing a compound having Formula II, such as Compound 2, or a composition comprising Compound 2, as described above. An amount of the compound(s) or composition(s) effective to inhibit viral replication is then administered to a
15 subject. The effective amount typically should be as high as the subject can tolerate. The currently preferred administration method is intravenous.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a graph of various concentrations of Compound 2 (μM) versus percent control
20 which illustrates the effects of Compound 2 on virus particles released from infected cells, where virus associated p24 antigen (\blacklozenge) was quantitated by antigen capture assay, RT activity (\blacksquare) was assessed by a homopolymeric(rA) template-primer system assay, and infectious units (\blacktriangle) were quantitated by titration of cell-free supernatant on MAGI cells.

FIG. 2 is a photograph of Western blot gels with AIDS patient serum or with polyclonal
25 antiserum to HIV-1 RT protein.

FIG. 3 is graph of concentration of Compound 2 versus percent control showing decreased (1) RT activity levels (\bullet), which were quantitated in the cell-free supernatant from TNF- α stimulated ACH2 cells in the presence of Compound 2, and (2) infectious units (\blacksquare), which were quantitated in the cell-free supernatant from TNF- α stimulated ACH2 cells in the
30 presence of Compound 2, (3) RT (\circ) of a separate sample, and (4) infectious units (\square) from a separate sample showing that under these conditions activities of RT and infectivity were recovered, where points on the graph represent means of triplicate tests from a representative

experiment. RT activity levels also were measured in virus harvested from drug-free TNF- α stimulated ACH2 cells after treatment of those preparations with either freshly prepared Compound 2 or with a fluid phase in which the virus had been cleared by centrifugation from the Compound 2 treated cultures.

5

DETAILED DESCRIPTION

I. Definitions


“Lower” as used herein refers to a compound or substituents having 10 or fewer carbon atoms in a chain, and includes all position, geometric and stereoisomers of such compounds or substituents.

“Aliphatic” refers to compounds having carbon and hydrogen molecules arranged in straight or branched chains including, without limitation, alkanes, alkenes and alkynes.

“Alkyl” as used herein refers generally to a monovalent hydrocarbon group formed by removing one hydrogen from an alkane. An alkyl group is designated generally as an “R” group, and has the general formula $-C_nH_{2n+1}$.

II. Compounds


Novel compounds of the disclosed invention have Formula I.



Formula I


With reference to Formula I, R is selected from the group consisting of hydrogen and lower aliphatic, particularly lower alkyl, such as methyl. Compound 2 is an example of a compound

5 having Formula I.




Compound 4


The present invention also is directed to a method of using compounds having Formula I and related biologically active compounds. These Formula I and related biologically active
5 compounds have Formula II.

**Formula II**

With reference to Formula II, R_1 is selected from the group consisting of hydrogen and lower aliphatic, particularly lower alkyl, such as methyl. R_2 is selected from the group consisting of –
5 CH_2COCH_3 and



Examples of such compounds include biologically active Compounds 1 (above) and 2 (below).

**Compound 2**

III. General Methods for Making BAIPs

Compound 2 can be made as described by Ryabova et al. in "2-Formyl-3-Aryl-aminoindoles in the Synthesis of 1,2- and 1,4-Dihydro-5H-Pyrido-[3,2-b]-Indole (δ Carboline) Derivatives," *Pharmaceutical Chemistry Journal*, 30:579-583 (1996), which is incorporated herein by reference. Other methods also can be used to make such compound, as well as other compounds according to the present invention. Example 1 describes a method for making Compound 2 as well.


With reference to Scheme 1, in a first such method IV was deacylated by action of Et_3N in methanol to form 3-p-nitrophenylaminoindole V, yield 80%, m.p. 220-222°C (MeOH), IR ν/cm^{-1} : 3350, 1590; MS m/z 253 (M^+). Formylation of V by treatment with Vilsmeier reagent produced the 2-formyl derivative VI, yield 96%, m.p. 237-238 °C (DMF- H_2O , 2:1); IR ν/cm^{-1} : 3290, 1640, 1600, 1575; ^1H NMR ($[\text{D}_6]\text{DMSO}$), δ : 9.88 (1H, s, CHO), 11.85, 9.41 (2H, 2s, NH, $\text{NHC}_6\text{H}_4\text{NO}_2$), 7.48 (4H, A_2B_2 system, $\text{C}_6\text{H}_4\text{NO}_2$), 6.95-7.59 (4H, m, arom. protons); MS m/z 281 (M^+).

Condensation of aldehyde VI with the dinitrile of malonic acid (both in the presence of Et_3N at 20°C or without Et_3N but under reflux) leads to dinitrile VII, yield 80% and 71%, respectively; a m.p. >270°C (dioxane); IR, ν/cm^{-1} : 3390, 3290, 2210, 1570; ^1H NMR ($[\text{D}_6]\text{DMSO}$), δ : 8.19 (1H, s, CH), 11.17, 9.68 (2H, 2s, NH, $\text{NHC}_6\text{H}_4\text{NO}_2$), 7.52 (4H, A_2B_2 system, $\text{C}_6\text{H}_4\text{NO}_2$), 7.11, 7.67 (4H, m, arom. protons); MS m/z 329 (M^+). Cyclization of

dinitrile VII can occur in either of two directions: with participation of *endo* (indole) or *exo* (at position 3) cyclic NH groups.

Heating VII in DMF-MeOH (1:1) caused intramolecular cyclization to form VIII isolated as the semihydrate (Scheme 1) yield 60%, m.p. 280 °C (decomp., DMF-MeOH, 1:1).

5



Scheme 1

Reagents and conditions for Scheme 1:

- i. Et_3N , MeOH, reflux two hours.
- ii. POCl_3 -DMF, 5-10°C, 0.25 hours; addition of a solution of 3 in DMF, standing of the mixture (20°C, 18 hours)
- 5 iii. PrOH , $\text{CH}_2(\text{CN})_2$, reflux 5 hours, or PrOH , PrOH , $\text{CH}_2(\text{CN})_2$, Et_3N , 5 hours, 20 °C.
- iv. DMF-MeOH, 1:1, reflux 0.25 hours.

Spectroscopic data for VII


IR ν/cm^{-1} : 3320, 2200, 1620, 1600, 1580.

- 10 ^1H NMR-DMSO- d_6 , S: 6.17(bs, 2H), 5.91 (d, 1H, H-C⁹), 6.74 (t, 1H, H-C⁸)
 ^{13}C NMR ($[\text{H}_6]$ DMSO) δ : 154.9 (C₂), 99.8 (C₃), 133.9 (C₄), 119.8 (C_{4a}), 114.5 (C_{9b}), 139.9 (C_{5a}), 128.8 (C_{9a}), 113.1, 119.9, 126.2, 127.1 (C₆₋₉), 119.9, 131.1 (C_{2,3,5,6}), 148.1, 144.1 (C_{1,4}), 117.7 (CN).

MS m/z 329 (M^+).

15

Scheme 2 shows an interesting and unexpected result that is obtained by methylating Compound VIII. Reacting VIII with methyl iodide in acetone in the presence of anhydrous K_2CO_3 adds the acetyl anion to the molecule's 4 position, together with tris-alkylation. As a result, 1-nitrophenyl-2-dimethylamino-3-cyano-4-acetyl-5-methyl-1,4-dihydropyrido[3,2-
20 b]indole X is obtained, yield 75%, m.p. 198-199 °C (MeOH-dioxane, 3:1).

**Scheme 2**

Reagent and conditions: MeI, acetone, anhydrous K_2CO_3 , reflux 56-60 hours, MeI added to the reaction mixture every 7-8 hours.

Spectroscopic data for X

IR ν/cm^{-1} : 1720, 2190.

$^1\text{H NMR}$ ($[\text{D}_6\text{H}_6]\text{DMSO}$) δ : 3.75 (3H, s, NMe-indole), 2.90 (6H, br.s, NMe_2), 2.10 (3H, s, CH_2COMe) 2.69 (2H, AB system $J_{\text{hem}}, 17$ Hz, $J^1_{\text{vic}} 9$ Hz, $J^2_{\text{vic}} 5$ Hz, CH_2COMe), 4.21 (1H, q, H- C_4), 7.89 (4H, A_2B_2 system, $\text{C}_6\text{H}_4\text{NO}_2$), 7.08-7.53 (4H, m, arom. protons). MS m/z 429 (M^+), 372 ($\text{M}^+ - \text{CH}_2\text{COMe}$).

In Scheme 2, first tris-methylation appears to occur with formation of a positively charged species, and the acetonide anion (formed in the reaction mixture in the presence of K_2CO_3) reacts at the electron-deficient position 4 to yield X. In the $^1\text{H NMR}$ spectrum of X (as distinct from VIII) a lower-field shift of the 9-H signal is not observed. The 1,4-dihydropyridine ring is not a flat system, and some data show that this ring has a boat conformation. Construction of molecular models for X, taking into account these data, shows that in this instance the p-nitrophenyl ring cannot influence the shape due to the anisotropic effect (as for VIII) and so the signals for all the protons in the condensed benzene ring are within the same range (7.08-7.53).


Ryabova et al., Khim.-Farm. Zh., 30: 42-45, (1996) reported the synthesis of 2-formyl-3-arylaminoindoles by formylation of the corresponding 3-arylaminoindoles according to the Vilsmeier reaction. Despite the "enamine" character of VI, the aldehyde group in position 2 is still capable of entering the reaction typical of this moiety. For example, reactions with primary amines lead to the formation of Schiff bases and the interactions with compounds possessing an active methylene group yield 2-vinylindole derivatives. Reaction of VI with malononitrile formed the 2-dicyanovinyl-3-arylaminoindoles VII, which are used to synthesize new indoles and condensed indole derivatives.

Heating compound VII for a short time in acetone in the presence of potassium carbonate leads predominantly to the hydration of vinyl fragment with the formation of initial aldehyde VI. The δ -carboline cyclization dominates when VII is heated in a DMF - MeOH (1:1) mixture up to the boiling temperature, and VIII is obtained at a 73% yield. The δ -carboline structure of VIII was confirmed by $^1\text{H NMR}$ spectroscopic data (Ryabova et al., Pharm. Chem. J. 30: 579-584, 1996). The $^1\text{H NMR}$ spectrum of VIII in $\text{DMSO}-d_6$ includes the following signals (δ , ppm): 6.17 (bs, 2H), 5.91 (d, 1H, H- C^9), 6.74 (t, 1H, H- C^8), 7.23 (t, 1H, H- C^7) and 7.42 (q, 1H, H- C^6).²⁾ 7.88 and 8.55 (A_2B_2 system, 4H, $\text{C}_6\text{H}_4\text{NO}_2$), 8.25 (s 1H, H- C^4). A characteristic feature of the latter spectrum is a considerable upfield shift of the H- C^9 proton

signal (5.91 ppm) as compared to the signals of other protons of the benzene ring (6.74 - 7.42 ppm) and the analogous proton signals in the spectra of pyrrolo[1,2-a]indole (7.27-7.94ppm) and 3-arylamino-2-formylindole X (6.95-7.59 ppm). Apparently, this shift of the H-C⁹ signal toward higher field strengths can be only due to the effect of anisotropic circular currents of the 4-nitrophenyl substituent in position 1, displaced out of the plane of the molecule as a result of steric constraints (the Dreiding molecular models). Thus, the experimental data confirmed the δ -carboline structure of VIII.

Alkylation of 3-aminoindole, initial aldehyde VI, 2-vinyl derivative VII, and 1,2-dihydro- δ -carboline VIII was used to develop a general method for making N-alkyl derivatives. This provided a common approach to obtaining compounds substituted at the exocyclic amino group and the nitrogen atom of the indole cycle. According to the mass-spectrometric data, methylation of VII by methyl iodide in acetone in the presence of potassium carbonate leads to the formation of a mixture of mono- and dimethyl derivatives, 2-formylindole, and δ -carboline X. Using column chromatography methods, aldehyde VI was isolated as was a bis-dimethyl derivative from this mixture. A side product in this reaction was 3-(4-nitrophenylamino)indole-2-carboxylic acid.

On heating in the presence of an aqueous alkali with dimethyl sulfate in acetone, compound VIII is methylated at the endo- and exocyclic nitrogen atoms (probably, via the stage of formation of the corresponding anion) yielding δ -carboline X from the reaction mixture (Scheme 3).



Scheme 3

The ^1H NMR spectrum of X (Table 2) contains signals from two methyl groups: $\delta = 3.18$ ppm (s, 3H, 2-NMe) and 3.81 ppm (s, 3H, 5-NMe). On saturation of the low-field N-methyl group signal, the intensity of the doublet at $\delta = 7.45$ ppm increases by 8%, and that of the singlet at $\delta = 8.50$ increases by 14%. In contrast, saturation of the signal of the other methyl group leads to no increase in the intensity of signals from aromatic protons. At the same time, saturation of the low-field part ($\delta = 7.70$ ppm) of the A_2B_2 system of signals from protons of the 4-nitrophenyl fragment increases by 4% the intensity of a doublet ($\delta = 5.82$ ppm) belonging to the proton at C^9 . The above NOE estimates unambiguously confirm the proposed structure of compound X, in which the methyl group at N^5 approaches the positions of $H-C^4$ and $H-C^6$, while the proton at C^9 is close to protons of the 4-nitrophenyl substituent in position 1. The comparatively small increase in intensity of the doublet due to C_9 protons ($\delta = 5.82$ ppm), observed on saturation of the signal from *ortho* protons of the nitrophenyl fragment, is probably explained by increasing distance to this proton system as a result of displacement of the N^1 -aryl substituent out of the molecular plane. This also leads to the upfield shift of the signal from $H-C^9$.

A different reaction of VIII with methyl iodide is observed in the presence of potassium carbonate, whereby the final result is determined by the methylation medium. For example, prolonged heating of the components in acetone leads to trimethylation of the initial carboline, accompanied by attachment of the acetonil anion in position 4. As a result, a tricyclic structure was obtained, in which the indole cycle is linked to the 1,4-dihydropyridine ring having a new functional substituent in position 4.

The dimethyl derivative X is apparently an intermediate involved in the formation of other compounds. This is confirmed by the fact that methylation of X using cyclohexanone or methylethylketone as solvents instead of acetone leads to 1-(4-nitrophenyl)2-dimethylamino-3-cyano-4-(2-oxocyclohexyl) and (3-oxo-2-butyl)-5-methyl-1,4-dihydro- δ -carbolines, respectively.

This initial stage may involve exhaustive methylation with the formation of a cation, in which the positive charge is delocalized between a dimethylamino group and position 4 of the molecule. It is this position to which the anion of a ketone (present in the reaction mass) is attached in the following stage with the formation of 1,4-dihydro- δ -carbolines.

The proposed structure of synthesized δ -carbolines was confirmed by spectroscopic data, primarily by the results of NMR measurements. For example, and with reference to

compound X, the IR spectrum of this compound, measured as a Nujol mull, showed the absorption bands at 1720 cm^{-1} (nonconjugated ketone) and 2190 cm^{-1} (CN group); mass spectrum (m/z): 429 $[M^+]$, 372 $[M^+ - \text{CH}_2\text{COCH}_3]$; ^1H NMR spectrum in DMSO- d_6 (δ , ppm): 3.75 (s, 3H, NMe), 2.90 (bs, 6H, NMe), 2.10 (s, 3H, CH_2COCH_3), 2.69 (AB-system, 2H, J_{hem} 17 Hz, J^1_{vic} 9 Hz, J^2_{vic} 5 Hz CH_2COCH_3), 4.31 (q, 1H, H-C⁴), 7.89 (A₂B₂-system, 4H, C₆H₄NO₂), 7.08 - 7.53 (4H, aromatic protons).

The IR spectra of synthesized compounds were measured on a Perkin-Elmer Model 457 spectrophotometer using samples prepared as Nujol mulls. The mass spectra were obtained on a Varian MAT-112 mass spectrometer with direct introduction of samples into the ion source operated at an ionizing electron energy of 70 eV. The NMR spectra were recorded on a Varian XL-200 instrument (USA) using TMS as the internal standard. The course of reactions was monitored and the samples were identified by thin-layer chromatography on Silufol UV-254 plates eluted in the chloroform methanol system (10:1). The data of elemental analyses coincided with the results of analytical calculations.

15

IV. Biological Activity

Compound 2 exerts broad anti-retroviral activity and has low cellular toxicity. Compound 2 initially was found active against HIV-1_{RF} in a standard screening cytoprotection assay ($\text{EC}_{50}=0.1\text{ }\mu\text{M}$ and a $\text{CC}_{50}>200\text{ }\mu\text{M}$) that requires multiple rounds of viral infection.

20 Range of action studies showed that Compound 2 also inhibited a panel of retroviruses, including laboratory and clinical isolates of HIV-1, HIV-1 isolates housing mutations that confer resistance to nucleoside and NNRTIs, monotropic and lymphotropic HIV-1 strains, as well as HIV-2 and SIV (Table 1).

Table 1 - Antiviral Properties of Compound 2

Virus	Cell	EC ₅₀	CC ₅₀	TI
HIV-1 RF	CEM-SS	0.1	>200	>2000
		0.078	>200	>2570
HIV-1 IIIB	CEM-SS	0.824	>200	>242
		0.836	126	151
HIV-1 OC/100	CEM-SS	4.68	116	24.9
		1.19	113	94.5
HIV-1 HEPT/236	CEM-SS	0.97	133	137
HIV-1 CALO-R	CEM-SS	1.10	122	110
		1.14	176	153
HIV-1 ddI-R	CEM-SS	0.62	163	263
HIV-1 DPS-R	CEM-SS	0.5	123	247
HIV-1 4X AZT	CEM-SS	1.3	110	84.8
HIV-1 A-17	CEM-SS	2.98	92.1	30.8
		3.48	88.1	25.3
HIV-1 6R/AZT	CEM-SS	16.6	130	7.8
		12.0	109	9.1
HIV-1 6S/AZT	CEM-SS	1.41	125	
		0.5	68.7	136
HIV-1 N119	CEM-SS	1.01	109	108
		9.73	124	12.8
HIV-2 ROD	CEM-SS	2.64	162	61.1
		4.79	>200	>41.7
		0.37	>200	>539
SIV	CEMx174	5.65	>200	>35.4
		6.5	134	20.6

- 5 ¹XTT antiviral assays were performed as described below in Example 4. EC₅₀ values indicate the drug concentration that provided 50% cytoprotection. CC₅₀ values reflect the drug concentrations that elicit 50% cell death. The XTT cytoprotection studies with HIV-1 were confirmed by measurement of supernatant RT, p24 and infectious virus titers.

- 10 Mechanistic studies showed no inhibitory activity of Compound 2 against RT when evaluated *in vitro* with recombinant p66/p51 RT using either the poly(rA) oligo(dT) or poly(rC) oligo(dG) template-primer systems. Likewise, Compound 2 did not affect virus binding or fusion to target cells, the activities of HIV-1 integrase or protease enzymes, or the nucleocapsid protein zinc fingers (Table 2).

Table 2 - Mechanism of Action Studies with Compound 2

Molecular Target ¹	Effect
RT (rAdT and rCdG)	NI ²
Protease	NI
Integrase	NI
NCp7 Zn fingers	NI
Biological Target	Effect
<u>Early Phase</u> HIV-1 Attachment Time Course Assay MAGI Assay	40% reduction at 100 μ M No inhibition of proviral DNA synthesis No reduction in blue cell formation at 200 μ M
<u>Late Phase</u> ACH-2 Assay	1) No reduction of p24 2) 2) Virus protein processing normal (Western blot) 3) Particle morphology normal (EM) 4) Reduction in RT activity in new virions 5) Reduction of infectious title of new virions

- ¹ Attachment of HIV-1 to CEM-SS cells, binding of gp120 to CD4, and the effects of compounds on HIV-1 RT, PR and NCp7 were quantitated as described below in Examples 4 and 5. ²NI indicates that no inhibition was observed at the high test concentration (100 μ M).

Thus, Compound 2 appeared not to act on any of the classical anti-HIV molecular targets.

- 10 The activity of Compound 2 was evaluated using a MAGI, cell-based, early-phase model of infection, described in Example 5. This assay requires virus binding, fusion, reverse transcription, integration of proviral DNA and the expression of Tat protein. Viruses were added to the MAGI cells in the presence or absence of Compound 2, and viral infectivity determined by scoring the number of blue foci. Compound 2 demonstrated no apparent
- 15 inhibitory action. Since the agent had no effect on these early-phase events, the data suggested it acted during the late phase of infection, after the HIV provirus integrates into the host cell genome.

Compound 2 was evaluated in a late-phase model of HIV-1 replication, described in Example 7. This model uses ACH2 cells, which carry a latent HIV-1 infection. In this model,

the ACH2 cells are treated with TNF- α which stimulates HIV-1 replication and virion production. Compound 2 had no effect on viral p24 antigen levels in the ACH2 cell culture supernatant, suggesting that virions were produced normally (FIG. 1). However, Compound 2 decreased virion-associated RT and viral infectivity levels in the culture supernatants in a concentration-dependent manner (FIG. 1). These observations were confirmed with latently infected U1 cells, chronically infected H9 cells, and other clones of latently infected ACH- 2 cells under TNF- α induced or uninduced conditions (data not shown).

With reference to FIG. 1, ACH2 cells were stimulated with recombinant TNF- α in the absence or presence of various concentrations of Compound 2. Cell-free supernatants were collected and evaluated as described in Examples 4-6. Virus-associated p24 antigen (◆) was quantitated by antigen capture assay, RT activity (■) was assessed by a homopolymeric(rA) template-primer system assay, and infectious units (▲) were quantitated by titration of the cell-free supernatant on MAGI cells wherein each blue cell represented an infectious unit. Examples 4-6. Each point represents the mean of triplicate cells from a representative experiment. Cell viability was unaffected at the relatively high test concentration of 200 μ M, as assessed by XTT assay.

The MAGI and ACH2 data, taken together, show that Compound 2 acts during the late phase of infection, after the provirus has integrated into the host cell genome. In the ACH2 assay, a drug which acted intracellularly to inhibit HIV replication would reduce the amount of HIV released into the cellular supernatant. However, HIV virions apparently being produced in an essentially normal manner, since Compound 2 treatment did not reduce the amount of viral p24 antigen present in the culture supernatant. However, when the HIV virions were released from the cell into the culture media, they exhibited significant abnormalities. Compound 2-treated cells showed reduced virion-associated RT activity and viral infectivity levels, and the degree to which the activity was reduced was directly related to the concentration of Compound 2.

To further investigate the observed abnormalities, the HIV-1 virions released from Compound 2-treated cells were compared to control in Western blot and protein analysis and electron microscopy. TNF- α stimulated ACH2 cells were treated with either Compound 2 or control solution, and cell-free supernatants were centrifuged to pellet the virus particles. Samples were subjected to Western blot analysis with AIDS patient serum or with polyclonal

antiserum to HIV-1 RT protein as shown by FIG. 2. The positions of gp120, Pr55^{gag} precursor polypeptide, p24 capsid (CA) protein, p17 matrix (MA) protein, integrase (IN), the p66 subunit of HIV-1 RT and p51 subunit of HIV-1 RT are indicated in FIG. 2. This analysis revealed a normal complement of fully mature (processed) HIV-1 proteins, including both subunits of the RT protein, in both control and Compound 2-treated supernatant. Electron micrographs of virus particles were obtained to assess morphological changes in virus particles treated with compounds of the present invention. Electron microscopy revealed no morphologic differences between virions obtained from control and Compound 2-treated cells. Thus, although virions released from Compound 2-treated cells had lower RT activity and were less infectious than virions released from control-treated cells, there were no abnormalities in virion morphology or protein composition that explained the difference.

A. Compound 2 is a Prodrug

The actual mechanism of action of Compound 2 became apparent partially from studies in which virion-associated RT levels were measured following centrifugation of virus particles in the virus-rich ACH-2 culture media. With reference to FIG. 3, RT activity (•) and infectious units (■) were quantified in the cell-free supernatant from TNF- α stimulated ACH2 cells in the presence of Compound 2. Activity levels decreased as the concentration of Compound 2 increased. A separate set of samples was centrifuged and the fluid phase removed prior to quantifying RT levels (○) and infectious units (□) of the virus pellet. Removing the culture fluid from the centrifuged virus particles allowed recovery of RT activities and virus infectivity at levels equivalent to those found in virions from untreated ACH-2 cultures (FIG. 3). This indicated that Compound 2 was a prodrug that had been converted into an active and reversible RT inhibitor during the 72-hour culture period. This was confirmed by a study in which the RT activity in a lysate of normal HIV-1 virions was inhibited by addition of virus-depleted culture supernatant from drug-treated ACH-2 cells. In contrast, addition of drug-free culture media or fresh drug to the normal virions did not inhibit their RT activity.

VI. Summary

Compounds 2 and 4 are novel RT inhibitors with truly broad-spectrum activity against retroviral RT enzymes and against infection by a broad range of retroviruses, including HIV-1, HIV-2 and SIV. BAIPs demonstrated antiviral activity against laboratory isolates of HIV-1 and

a panel of clade-representative clinical isolates in PBMC cultures at submicromolar levels. More impressive though was the ability of the BAIPs to inhibit the replication of a panel of HIV-1 variants carrying mutations in RT that confer resistance to AZT and various NNRTIs such as oxithiin carboxanilide (L-100→I), thaizolobenzimidazole (V-108→I), calanolide (T-139→I), diphenylsulfone (Y-181→I), 3TC (M-184→I) and others. The ability of the BAIPs to inhibit the enzymatic RT activities and replication of this wide array of retroviruses distinguished it from classical NNRTI type molecules that are HIV-1 specific and can be typically rendered ineffective by one or more single mutations in the HIV-1 RT enzyme. Thus, the BAIPs truly represent the first reported example of a broadly antiretroviral NNRTI (BANNRTI).

The BAIPs have been found to inhibit not only all strains of HIV-1 tested, but also the replication of HIV-2 and SIV. This property sets the BAIPs apart from other NNRTI-type agents. The BAIPs may be used for therapy to individuals already carrying HIV-1 variants that are resistant to AZT or classical NNRTI molecules.

Classical NNRTIs bind noncovalently to the non-substrate binding site of the RT enzyme, and mutations in this region of the enzyme result in loss of sensitivity to the agents. Likewise, nucleoside analogs interact with RT in the substrate binding pocket, and mutations in this region of the enzyme result in resistance to the respective nucleoside analogs. Because BAIPs exert such distinct antiviral properties from the classical NNRTIs and have such a different structure from nucleoside analogs, BAIPs likely interact with RT in a different manner that classical NNRTIs. A series of computational studies were performed that predict the most likely binding site for BAIPs. Such studies suggested that BAIPs bound tightly in a previously unidentified pocket near the Asp triad in the active site of the RT enzyme. Together, these studies set the BAIP molecules apart as a new class of RT inhibitors, the BANNRTIs.

25

VI. Pharmaceutical Compositions Comprising Compounds 1 and 2

The vehicle in which disclosed compounds can be delivered include pharmaceutically acceptable compositions of the drugs. Any of the common carriers, such as sterile saline or glucose solution, can be used with the compounds provided by the invention. Routes of administration include, but are not limited to, oral and parenteral routes, such as intravenous (iv), intraperitoneal (ip), rectal, topical, ophthalmic, nasal, transdermal, and combinations thereof.

The drugs may be administered intravenously in any conventional medium for intravenous injection, such as an aqueous saline medium, or in blood plasma medium. The medium also may contain conventional pharmaceutical adjunct materials such as, for example, pharmaceutically acceptable salts to adjust the osmotic pressure, lipid carriers such as cyclodextrins, proteins such as serum albumin, hydrophilic agents such as methyl cellulose, detergents, buffers, preservatives and the like. A more complete explanation of parenteral pharmaceutical carriers can be found in *Remington: The Science and Practice of Pharmacy* (19th Edition, 1995) in chapter 95. The compositions are preferably in the form of a unit dose in solid, semi-solid and liquid dosage forms such as tablets, pills, powders, liquid solutions or suspensions.

VII. Administering Compounds

The present invention provides a treatment for HIV and SIV disease, perhaps by RT inhibition, and associated diseases, in a subject such as an animal, for example a monkey or human. The method includes administering a compound, or compounds, of the present invention, or a combination of the compound or compounds and one or more other pharmaceutical agents. The compound, or compounds, can be administered to the subject in a pharmaceutically compatible carrier. The compound, or compounds, are administered in amounts effective to inhibit the development or progression of HIV and SIV disease. Although the treatment can be used prophylactically in any patient at significant risk for such diseases, subjects can also be selected using more specific criteria, such as a definitive diagnosis of the condition.

The disclosed compounds are ideally administered as soon as possible after potential or actual exposure to viral infection. For example, once viral infection has been confirmed by laboratory tests, a therapeutically effective amount of the drug is administered. The dose can be given by frequent bolus administration.

Therapeutically effective doses of the compounds of the present invention can be determined by one of ordinary skill in the art. For example, effective doses can be such as to achieve tissue concentrations that are at least as high as the EC₅₀. The low cytotoxicity of the BAIP makes it possible to administer high doses, for example 100 mg/kg, although doses of 10 mg/kg, 20 mg/kg, 30 mg/kg or more are contemplated. Thus, the dosage range likely is from about 0.1 to about 200 mg/kg body weight orally in single or divided doses, more likely from

about 1.0 to 100 mg/kg body weight orally in single or divided doses. For oral administration, the compositions are, for example, provided in the form of a tablet containing from about 1.0 to about 1000 mg of the active ingredient. Symptomatic adjustment of the dosage to the subject being treated can be achieved by suing tablets of varying amounts of compound, such as 1, 5,
5 10, 15, 20, 25, 50, 100, 200, 400, 500, 600, and 1000 mgs of the active ingredient.

The specific dose level and frequency of dosage for any particular subject may be varied and will depend upon a variety of factors as will be known to a person of ordinary skill in the art. These include the activity of the specific compound, the metabolic stability and length of action of that compound, the age, body weight, general health, sex, diet, mode and time of
10 administration, rate of excretion, drug combination, and severity of the condition of the host undergoing therapy.

The pharmaceutical compositions can be used in the treatment of a variety of retroviral diseases caused by infection with retroviruses that require reverse transcriptase activity for infection and viral replication. Examples of such diseases include HIV-1, HIV-2, and the
15 simian immunodeficiency virus (SIV).

The present invention also includes combinations of a BAIP compound, or BAIPs, of the present invention with one or more agents useful in the treatment of viral diseases, such as HIV disease. For example, the compounds of this invention may be administered, whether before or after exposure to the virus, in combination with effective doses of other antivirals,
20 immunomodulators, anti-infectives, or vaccines. The term "administration" refers to both concurrent and sequential administration of the active agents.

Examples of antivirals that can be used in combination with the BAIP RT inhibitors of the invention are: AL-721 (from Ethigen of Los Angeles, CA), recombinant human interferon beta (from Triton Biosciences of Alameda, CA), Acemannan (from Carrington Labs of Irving,
25 TX), ganciclovir (from Syntex of Palo Alto, CA), didehydrodeoxythymidine or d4T (from Bristol-Myers-Squibb), EL10 (from Elan Corp. of Gainesville, GA), dideoxycytidine or ddC (from Hoffman-LaRoche), Novapren (from Novaferon labs, Inc. of Akron, OH), zidovudine or AZT (from Burroughs Wellcome), didanosine, lamiduvine, delavirdine, nevirapine, ribavirin (from Viratek of Costa Mesa, CA), alpha interferon and acyclovir (from Burroughs Wellcome),
30 indinavir (from Merck & Co.), 3TC (from Glaxo Wellcome), Ritonavir (from Abbott), Saquinavir (from Hoffmann-LaRoche), nelfinavir, and others.

Examples of immunomodulators that can be used in combination with the BAIPs of the invention are AS-101 (Wyeth-Ayerst Labs.), bropiramine (Upjohn), gamma interferon (Genentech), GM-CSF (Genetics Institute), IL-2 (Cetus or Hoffman-LaRoche), human immune globulin (Cutter Biological), IMREG (from Imreg of New Orleans, La.), SK&F106528, and
5 TNF (Genentech).

Examples of some anti-infectives with which the BAIPs can be used include clindamycin with primaquine (from Upjohn, for the treatment of pneumocystis pneumonia), fluconazole (from Pfizer for the treatment of cryptococcal meningitis or candidiasis), nystatin, pentamidine, trimethaprim-sulfamethoxazole, and many others.

10 The combination therapies are not limited to the lists provided, but include any composition for the treatment of HIV disease and related retroviral diseases (including treatment of AIDS).

VI. EXAMPLES

15 The following examples are provided to exemplify certain particular features of working embodiments of the present invention. The scope of the present invention should not be limited to those features exemplified.

Example 1

20 This example describes methods for making Compound 2 and related compounds.

2-Cyano-3-[3-(4-nitrophenylamino)-2-indolyl]acrylic acid nitrile (VII, Scheme 1).

Method 1. A mixture of 3.65 g (13 mmole) of compound VI, 1.6 g (24 mmole) malononitrile, 0.25 ml (2 mmole) triethylamine, and 73 ml of 2-propanol was stirred for 5 h at 20°C and allowed to stand at this temperature for 16 h. The precipitate was separated by
25 filtration and washed with 2-propanol to obtain 3.3 g of VII.

Method 2. A mixture of 3 g (11 mmole) of Compound VI, 1.5 g (22 mmole) malononitrile, and 60 ml of 2-propanol was refluxed for 4 h and allowed to stand for 16 h at 20°C. Then the reaction mixture was treated as in method 1 to obtain 2.7 g of VII.

Method 3. A suspension of 0.3 g (1 mmole) of N-acetylated derivative of VI, 0.1 g (1.5
30 mmole) malononitrile, and 0.13 g (1.5 mmole) fused sodium acetate in 5 ml of acetic acid was stirred for 0.5 h at 20°C, followed by 3 h at 80°C. Then 0.1 g of malononitrile was added and

the mixture was stirred for another 5 h at 20°C. Then the mixture was cooled, and the precipitate was separated by filtering and washed with AcOH, water, and MeOH to obtain 0.05 g of VII.

1-(4-Nitrophenyl)-2-imino-3-cyano-1,2-dihydro-5H-pyrido[3,2-b]indole (VIII, Scheme 1).

Method 1. A mixture of 3.3 g (10 mmole) of nitrile VII, 15 ml MeOH, and 15 ml DMF was heated to boiling. As a result, VII dissolved and a new precipitate appeared. This suspension was refluxed for 5 min and cooled. The precipitate was separated by filtering and washed with MeOH to obtain 2.4 g of VIII. ¹³C NMR spectrum in DMSO-d₆ (δ, ppm): 154.9 (C²), 99.8 (C³), 133.9 (C⁴), 119.8 (C^{4a}), 114.5 (C^{9b}), 139.9 (C^{5a}), 128.8 (C^{9b}), 113.1, 119.9, 126.2, 127.1 (C⁶-C⁹), 119.9, 131.1 (C^{2'}, C^{3'}, C^{5'}, C^{6'}), 148.1, 144.1 (C^{1'} C^{4'}), 117.7 (CN).

Method 2. A mixture of 0.33 g (1 mmole) of nitrile VII and 0.4 g (3 mmole) of calcined potassium carbonate in 10 ml of acetone was refluxed for 15 min. The precipitate was separated by filtering and washed with water to obtain 0.05 g of VIII. The acetone mother liquor was evaporated, and the residue triturated with diethyl ether to obtain 0.17 g (61%) of VIII.

Methylation of 3-(4-nitrophenylamino)indole. To a mixture of 1.3 g (5 mmole) of 3-(4-nitrophenylamino)indole, 16 ml DMF, and 2.1 g (15 mmole) of calcined potassium carbonate was added 5 ml MeI and the mixture was stirred at 80°C for 60 h, with 2 ml MeI added each 6 h (to a total of 20 ml). The mixture was cooled, the remaining potash separated by filtering and washed with DMF, and the filtrate was evaporated. The residue was triturated with diethyl ether on adding a minimum amount of MeOH and filtered. The filtrate was evaporated, and the residue chromatographed on a silica gel column with chloroform. Five sequential 100 ml fractions were collected, and the third and fifth fractions containing individual products were evaporated. Fraction 1 yielded 0.6 g (42%) of 1-methyl-3-[N-methyl-N-(4-nitrophenyl)amino]indole, and fraction 3 yielded 0.4 g of 3-[N-methyl-N-(4-nitrophenyl)amino]indole.

1-(4-Nitrophenyl)-2-methylimino-3-cyano-5-methyl-1,2-dihydro-5H-pyrido[3,2-b]indole (XIV, Scheme 3). To a solution of 2 g (50 mmole) of NaOH in 2 ml water was added 100 ml acetone and 3.3 g (10 mmole) of VIII, and the mixture was heated to boiling on stirring and refluxed for 5 min. To this mixture was added 4 ml (40 mmole) of Me₂SO₄ and the boiling

was continued with stirring for 6 h. Another 4 ml of Me_2SO_4 was added and the mixture was refluxed for another 6 h. Then the mixture was cooled, the precipitate separated by filtration, washed with acetone, and dissolved in 500 ml of boiling water. The solution was filtered hot, cooled and alkalified with 1N KOH (15 ml). The precipitate was filtered and washed
5 sequentially with water, 2-propanol, and diethyl ether to obtain 2.1 g of XIV.

1-(4-Nitrophenyl)-2-dimethylamino-3-cyano-4-(2-oxo-propyl)-5-methyl-1,4-dihydro-5H-pyrido-[3,2-b]indole (XI, Scheme 3)

Method 1 . To a suspension of 2.15 g (6.5 mmole) of VIII and 3.6 g (26 mmole) of calcined potassium carbonate in 80 ml of acetone was added 2 ml MeI and the mixture was
10 refluxed on stirring for 60 h, with 2 ml MeI added each 7 - 8 h. Then the mixture was cooled and the remaining potash separated by filtering and washed with acetone. The filtrate was evaporated, and the residue triturated with water, filtered, and washed with water and methanol to obtain 2.1 g of a technical-purity product 1-(4-nitrophenyl)2-dimethylamino-3-cyano-4-(oxo-propyl)-5-methyl-1,4-dihydro-5H-pyrido[3,2-b]indole. The product was purified by boiling with
15 20 ml MeOH, after which the insoluble precipitate was filtered to obtain 1.5 g of Compound 2-(4-nitrophenyl)2-dimethylamino-3-cyano-4-(oxo-propyl)-5-methyl-1,4-dihydro-5H-pyrido[3,2-b]indole.

Method 2. A mixture of 1.07 g (3 mmole) of XIV, 0.83 g (6 mmole) calcined potassium carbonate, 70 ml acetone, and 2 ml MeI was refluxed with stirring for 45 h, followed
20 by a procedure similar to that in method 1. This yielded 0.85 g of X, which was identical to the product obtained by method 1.

1-(4-Nitrophenyl)-2-dimethylamino-3-cyano-4-(2-oxo-2-butyl)-5-methyl-1,4-dihydro-5H-pyrido-[3,2-b]indole. To a suspension of 0.33 g (1 mmole) of VIII and 0.65 g (4.7 mmole) of calcined potassium carbonate in 20 ml of methylethylketone was added 2 ml
25 MeI . The mixture was refluxed with stirring for 41 h, with 2 ml MeI added each 6 h. The mixture was cooled and the remaining potash separated by filtering and washed with diethyl ether, water, and methanol. The residue was mixed with chloroform and the solution filtered and evaporated. The residue was triturated with ether, and the precipitate was filtered and washed with ether to obtain 0.1 g of 1-(4-nitrophenyl)-2-dimethylamino-3-cyano-4-(2-oxo-2-
30 butyl)-5-methyl-1,4-dihydro-5H-pyrido-[3,2-b]indole.

Example 2

This example describes virus replication inhibition assays that have been performed. The established human cell lines and laboratory-derived virus isolates (including drug resistant virus isolates) used in these evaluations have previously been described (Weislow et al., 1989; Rice and Bader, 1995). The antiviral activities and toxicity profiles of the compounds were evaluated with CEM-SS cells and HIV-1_{RF} using the XTT (2,3-bis[2-methoxy-4-nitro-5-sulfophenyl]-5-[(phenylamino)carbonyl]-2H-tetrazolium hydroxide) cytoprotection microliter assay which quantifies the ability of a compound to inhibit virus-induced cell killing or to reduce cell viability itself (Weislow et al., 1989; Rice and Bader, 1995). The data are reported as the concentration of drug required to inhibit 50% of virus-induced cell killing (EC₅₀) and the concentration of drug required to reduce cell viability by 50% (CC₅₀). HIV-1 isolates utilized included common laboratory strains (RF, IIB and MN), as well as a panel of HIV-1 clinical isolates (Rice et al., 1997). The pyridinone-resistant HIV-1_{A17} isolate was obtained from Emilio Emini at Merck Sharpe and Dohme Laboratories. CEM, U1, ACH-2, HeLa-CD4-LTR- β -gal, 174xCEM, and H9/HTLV-III_B NIH 1983 cell lines were obtained from the AIDS Research and Reference Reagent Program (National Institute of Allergy and Infectious Disease, National Institutes of Health, Bethesda, MD), as were the HIV-2_{ROD} and the SIV isolates. Phytohemagglutinin-stimulated human peripheral blood lymphocytes and monocyte/macrophages were prepared and utilized in antiviral assays as previously described (Rice et al., 1996), and levels of virion-associated p24 in cell-free culture supernatants were determined via antigen capture ELISA (Beckman Coulter).

Example 3

This example describes integrase, protease, RT and NC zinc finger assays that have been performed. *In vitro* inhibitory activity against recombinant HIV-1 protease was performed with a reverse-phase high-pressure liquid chromatography assay utilizing the Ala-Ser-Glu-Asn-Tyr-Pro-Ile-Val-Glu-amide substrate (multiple Peptide System, San Diego, CA.) (Rice et al., 1993a). The *in vitro* actions of compounds on 3'-processing and strand transfer activities of recombinant HIV-1 integrase were assayed according to Bushman and Craigie (1991), but with modifications (Turpin et al., 1998). The action of compounds on the RNA-dependent polymerase activity of recombinant HIV-1 p66/p51 RT was determined by measuring incorporation of [³²P]TTP or [³²P]GTP into the poly rA:oligo dT(rAdT) or poly rC:oligo

dG(rCdG) homopolymer template-primer systems, respectively, while the inhibition of drug on the DNA-dependent polymerase activity of purified recombinant HIV-1 RT was determined by measurement of incorporation of [32 P]TTP or [32 P]GTP into the polydA:oligo(dT)dAdT or polydC:oligo(dG)dCdG homopolymer template-primer systems, respectively (Pharmacia Biotech, Piscataway, NJ). Reactions were performed in the presence or absence of the drug as described previously (Rice et al., 1997). Reactions were terminated with ice-cold 10% trichloroacetate, filtered through GF/C filter under vacuum, and the filters were then washed with 100% ethanol and [32 P] incorporation quantitated by Cerenkov counter. The LTR region of the HIV-1 genomic RNA was prepared from a pGEM LTR by in vitro transcription with T7 RNA polymerase (Promega, Madison, WI). In pGEM LTR, LTR region from pNL4-3 was inserted into the polylinker of pGEM (Promega) in the orientation that the sense LTR RNA were made when T7 RNA polymerase was used. The rest of steps for the preparation of heteropolymeric primer-template and RT reaction was performed as described (Gu et al., 1993).

Virion-associated RT activity was performed as described previously (REF) in the presence or absence of compound with the homopolymeric template-primer (rAdT, rCdG, dAdT and dCdG) (Pharmacia Biotech, Piscataway, NJ) or heteropolymeric template-primer prepared as described above. HIV-2_{ROD10} and SIV virions were obtained by transfection of proviral DNA into HeLa cells.

20

Example 4

This example describes RNase H cleavage assays that have been performed. An α -[32 P]-uridine-labeled RNA template (81 nucleotides in length) was hybridized to a 20-base DNA oligonucleotide in the presence of 50 mM Tris-HCl, pH 8.0, 50 mM NaCl, 2.0 mM dithiothreitol, 100 μ g/ml acetylated bovine serum albumin, and 10 mM CHAPS as previously described (Gao et al., 1998). For these reactions, 100 ng of RNA (approximately 50,000 cpm) and 20 ng of DNA (oligonucleotide 3352, 5'TTCTCGACCCTTCCAGTCCC 3') were utilized. Purified HIV-1 wild type RT (45 ng) was mixed with COMPOUND 4 such that the final concentrations were 0.1, 1.0, 10 or 100 μ M, and the reactions were initiated by the addition of 60 mM MgCl₂ and the annealed RNA/DNA complex in a final volume of 12 μ l. This mixture was incubated at 37°C for 1 minute with Compound 4 or for various times without the compound. Reactions were terminated by the addition of 2X loading buffer, and the products were heat denatured and resolved on a 15% denaturing polyacrylamide-7M Urea gel in TBE

buffer at 1600 Volts for approximately 90 minutes. Gels were dried and exposed for autoradiography overnight, and the film was developed with a Kodak RP X-OMAT processor.

Example 5

5 This example describes MAGI cell assays that have been performed. The MAGI cell indicator line was obtained from the AIDS Research and Reference Program, Division of AIDS, National Institute of Allergy and Infectious Disease. MAGI cells are a HeLa cell line that both expresses high levels of CD4 and contains a single integrated copy of a beta-galactosidase gene under the control of a truncated human immunodeficiency virus type 1 (HIV-1) long terminal repeat (LTR). These cells maintained in DMEM medium supplemented with 5% fetal bovine serum (FBS), 100U of penicillin G sodium, 0.1 mg of streptomycin sulfate, 0.2 mg G418 sulfate, and 0.1 mg of hygromycin B per ml.

MAGI cells and an HIV-1 *env*- and *Tat*-expressing HeLa (HL2/3) cell line were used to perform a fusion assay. *Tat* activates gene expression from the HIV LTR, and therefore upon
15 fusion of MAGI and HL2/3 cells, *tat* expressed in HL2/3 cells (Ciminale et al., 1990) would activate β -galactosidase expression in MAGI cells. MAGI or HL2/3 cells (2.5×10^5 in 500 μ l 5%FBS/DMEM) were preincubated with the tested compound for 1 hour at 37° C, respectively. At the end of preincubation, two cell lines were mixed at 1:1 ratio and were continued incubated for another 16 hours. The cells were then fixed and stained for the expression of β -
20 galactosidase with indolyl- β -D-galactopyranoside (X-Gal) as described previously (Kimpton and Emerman, 1992). The numbers of blue cells were counted by light microscopy.

MAGI cells were also used to examine the effects of compounds on virus replication, from attachment through early gene expression. In these assays, the LTR-driven β -galactosidase gene in MAGI cells would not be activated until the incoming virus had penetrated the cell,
25 reverse transcribed its RNA genome, generated the double-stranded proviral DNA, integrated the proviral DNA into the host cell genome, and expressed its *tat* gene. The assay was preformed as previously described with modifications (Howard et al., 1998). The virus stock used in the assay was prepared either from TNF- α -induced U1 cells (HIV_{IIIb}) or pNL4-3 - transfected from HeLa cells transfected with the pNL4-3 plasmid containing HIV-1 proviral
30 DNA. Viruses were diluted in 200 μ l DMEM medium supplemented with 5% fetal bovine serum (FBS), and were titrated to generate approximately 300 blue cells per well in 24 well plates. Viruses were added to the MAGI cells in the presence or absence of the test compound.

After 2 hours incubation at 37° C, the virus was removed, the cells were washed and 1 ml 5%FBS/DMEM medium with or without the test compound was added to the cells. For the time-of-addition assay, the compound was added at time zero when the infection was initiated, or at 2, 4, 8 or 24 hours post initiation of the infection. For the time-of-removal assay, the compound was added to all wells at the beginning of infection and was then removed at 2, 4, 8, 24 or 48 hours thereafter. The cells were washed once with medium after removal of the drug followed by the readdition of 1ml 5%FBS/DMEM fresh medium. Forty-eight hours post initiation of infection, cells were fixed and stained as described above.

To titrate the infectivity of viruses harvested from the drug-treated chronic infected cells, MAGI cells were also used. Either 500 µl total culture media or 200 µl pelleted viruses were added to the 24 well culture plates in the presence 20 µg/ml DEAE-dextran for 3 hours at 37° C prior to the addition of 2 ml of media. The cultures were fixed and stained as described above.

Example 6

This example describes PCR analysis of nascent proviral DNA. MAGI cells were plated at a density of 4×10^5 /well in a 6-well plate. Twenty-four hours later, the cells were infected with HIV_{IIIB} viruses in 500 µl 5%FBS/DMEM in the presence or absence of the compound. HIV_{IIIB} viruses were prepared from TNF-α-induced U1 cells and the amount used in one infection was titrated as the amount producing 1000 blue colonies. Four hours post-infection, the cells were trypsinized, washed and digested at 55° C for 1 hour with 100 µg/ml protease K in 100 µl buffer containing 0.5% Triton X-100, 100 mM NaCl, 50 mM Tris (pH 7.4), and 1 mM EDTA. To inactivate protease K, the samples were then heated at 100° C for 15 minutes. PCR reactions were performed using M661 and M667 primers (Zack et al., 1990) and 5 µl sample was used in each reaction.

Example 7

This example describes ACH2 latently-infected cell assays that have been performed. ACH2 cells were maintained in RPMI 1640-10% FBS medium. Forty thousand ACH2 cells per milliliter were induced with 5 ng of recombinant tumor necrosis factor alpha (TNF-α) (Sigma Chemical Co., St. Louis, MO) per ml for 24 hours. Twenty-four hours later, an equal volume of medium supplemented with 5 ng of TNF-α per ml and with the appropriate (2x final)

concentration of the tested compound was added to cells. Viruses containing cell-free supernatants were collected 48 hours later, and they were subjected directly or after being pelleted through centrifugation for RT assay, p24 assay, and virus titration assay. Viability of the cultures was determined by XTT dye reduction). The RT assay, virus titration assay with
5 MAGI cells, and p24 assay were performed as described above.

Pelleted virus particles were also subjected to Western blot analysis. The virion-associated viral proteins pelleted from 400 μ l of cell free supernatant were resolved on 10% SDS-polyacrylamide gels, were electroblotted onto polyvinylidene difluoride (PVDF) membranes, and were detected by AIDS patient sera or by rabbit-polyclonal anti-HIV-1 RT
10 antibody (AIDS Research and Reference Program, Division of AIDS, National Institute of Allergy and Infectious Disease). Western blots were developed with standard methodology by chemiluminescence (Dupont-NEN, Wilmington, Del.) with a goat-anti human or goat anti-rabbit horseradish peroxidase-conjugated antibody (Bio-Rad, Hercules, Calif.).

15 **Example 8**

This example describes molecular modeling that has been done concerning BAIPs. The following analysis was carried out on the HIV-1 RT coordinates 1RTH (Abola et al., 1987; Bernstein et al., 1977). A two-stage analysis was performed. First, the exterior surface of the HIV-1 RT heterodimer was probed for candidate binding regions. This process consists of
20 localized sampling of the solvent accessible surface to determine a statistical probability that a candidate ligand may bind at this site. The model used to make the calculation has been parameterized, based on a broad sampling of protein-ligand crystal complexes available in the Brookhaven database of protein structures. (PDB) (Abola et al., 1987; Bernstein et al., 1977). The complete details for identification of putative protein binding sites can be found in Young
25 et al. (Young et al., 1994). Second, the optimal docked position of the test ligand was determined. Families of possible conformations for the test ligand were generated using standard modeling techniques and each was docked to the regions defined in the first step. The docking procedure has been demonstrated to have an accuracy of within 1Årms deviation from the known docked positions (Wallqvist & Covell, 1996). The position of the ligand with the
30 strongest calculated binding strength is reported herein.

Example 9

This example describes the preparation of samples for electron microscopy. Sample preparation for electron microscopy is described previously (Gonda et al., 1985). Briefly, the virus pellets were fixed with a 0.1M sodium cacodylate buffer containing 1.25% glutaraldehyde, pH 7.2, followed by a 1% osmium tetroxide in the same buffer. The fixed pellets were dehydrated in a series of graded ethanol solutions (35%, 50%, 75%, 95% and 100%) and propylene oxide. The pellets were infiltrated overnight in an epoxy resin (LX-112) and propylene oxide mixture, then embedded in epoxy resin to cured for 48 hours at 60C. Thin-sections (50 to 60 nm) of the pellet were cut, mounted on a naked copper grid, and double stained with uranyl acetate and lead citrate. The thin sections were stabilized by carbon evaporation in a vacuum evaporator, observed, and photographed with a Hitachi H-7000 electron microscope operated at 75kv.

The present invention has been described with respect to certain embodiments. The scope of the invention should not be limited to these described embodiments, but rather should be determined by reference to the claims.

References

1. Abola, E.E., et al. (1987). In *Crystallographic Databases – Information Content, Software Systems, Scientific Applications*, (Allen, F. H., Bergerhoff, G. & Sievers, R., eds), pp. 107-132.
2. Bernstein, F.C., et al. (1997) *J. Mol. Biol.* 112:535-542.
3. Baba, M., et al. (1989) *Biochem. Biophys. Res. Comm.* 165:1375-1381.
4. Balzarini, J. et al. (1992) *Proc. Natl Acad. Sci USA.* 89:4392-4396.
5. Buckheit, R.W. et al. (1997) *Antimicrob. Agents Chemother.* 41:831-837.
- 25 6. Bushman, F.D., and R. Craigie (1991) *Proc. Natl. Acad. Sci. USA* 88:1339-1343.
7. Ciminale, V. et al. (1990) *AIDS Res. Hum. Retrovir.* 6:1281-1287.
8. Cohen, J. (1997) *Science* 277:32-33.
9. Cohen, K. A. et al. (1991) *J. Biol. Chem.* 266:14670-14674.
10. Condra, J. H. et al. (1992) *Antimicrob. Agents Chemother.* 36:1441-1446.
- 30 11. De Clercq, E. (1992) *AIDS Res. Hum. Retrovir.* 8:119-134.
12. De Clercq, E. (1996) *Rev. Med. Virol.* 6:97-117.
13. Debyser, Z. et al. (1991) *Proc. Natl. Acad. Sci. USA.* 88:1451-1455.

14. Finzi, D et al. (1997) *Science* 278:1295-1300.
15. Gao, H.Q., et al. (1998) *J. Mol. Biol.* 277:559-572.
16. Goldman, M. E. et al. (1991) *Proc. Natl. Acad. Sci. USA.* 88:6863-6867.
17. Goldman, M. E., et al. (1992) *Antimicrob. Agents Chemother.* 36:1019-1023.
- 5 18. Gonda, M.A., et al. (1985) *Science* 227:173-177.
19. Gu et al. (1993) *J. Biol Chem* 269:28119-28122.
20. Gao et al. (1998) *J. Mol. Biol.* 1998, 277: 559-572
21. Howard, O.M.Z. et al. (1998) *J. Med. Chem.* 41:2184-2193.
22. Kimpton, J., and M. Emerman (1992) *J. Virol.* 66:2232-2239.
- 10 23. Koup, R. A. et al. (1991) *J. Infect. Dis.* 163:966-970.
24. Mellors, J.W. (1995) *Intl. Antiviral News* 3:8-13.
25. Merluzzi, V. J. et al. (1990) *Science* 250:1411-1413.
26. Miyasaka, T., H. et al. (1989) *J. Med. Chem.* 32:2507-2509.
27. Pauwels, R., K. et al. (1993) *Proc. Natl. Acad. Sci. USA.* 90:1711-1715.
- 15 28. Pauwels, R., K. et al. (1990) *Nature* 343:470-474.
29. Quinones-Mateu, M.E. et al. (1997) *J. Virol.* 29:364-373.
30. Rice, W.G. et al. (1997) *Antimicrob. Agents Chemother.* 41:419-426.
31. Rice, W.G., et al. (1993) *Nature* 361:473-475.
32. Rice, W.G. et al. (1995) *Science* 270:1194-1197.
- 20 33. Rice, W.G. et al. (1996) *J. Med. Chem.* 39:3606-3616.
34. Rice, W.G. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:9721-9724.
35. Rice, W.G. and J.P. Bader (1995) in *Advances in Pharmacology* (J. T. August, M. W. Anders, F. Murad and J. T. Coyle, eds) pp. 389-438.
36. Romero, D. L. et al. (1991) *Proc. Natl. Acad. Sci. USA.* 88:8806-8810.
- 25 37. Romero, D. L et al. (1993) *J. Med. Chem.* 36:1505-1508.
38. Ryabova et al.(1996) *Pharm. Chem. J.* 30:579-584.
39. Ryabova et al.(1996) *Khim.-Farm. Zh.*30:42-45
40. Ryabova et al. (1996) *Mendeleev Commun.* 3:107-109
41. Turpin, J.A. (1997) *Antiviral Chem Chemother.* 8:60-69.
- 30 42. Wallqvist, A. and D.G. Covell (1996) *Proteins* 25:403-419.
43. Weislow, O.S. (1989) *J. Natl. Cancer Inst.* 81:577-586.
44. Wishka, D.G. et al. (1998) *J. Med. Chem.* 41:1357-1360.

45. Wong, J.K. et al. (1997) *Science* 278:1291-1295.
46. Young, L. et al. (1994) *Prot. Sci.* 3:717-729.
47. Zack, J.A., et al. (1990) *Cell* 61:213-222.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2